arXiv:1210.4225v1 [hep-ph] 16 Oct 2012

Neutrinos from Pion Decay at Rest to Probe the Proton Strangeness

G. Pagliaroli,[1, *] C. Lujan-Peschard,[1, 2, †] M. Mitra,[1, ‡] and F. Vissani[1, §]
[1] *INFN, Laboratori Nazionali del Gran Sasso, Assergi (AQ), Italy*
[2] *Departamento de Fisica, DCeI, Universidad de Guanajuato, León, Guanajuato, México*

The study of the neutral current elastic scattering of neutrinos on protons at lower energies can be used as a compelling probe to improve our knowledge of the strangeness of the proton. We consider a neutrino beam generated from pion decay at rest, as provided by a cyclotron or a spallation neutron source and a 1 kton scintillating detector with a potential similar to the Borexino detector. Despite several backgrounds from solar and radioactive sources it is possible to estimate two optimal energy windows for the analysis, one between $0.65-1.1$ MeV and another between $1.73-2.5$ MeV. The expected number of neutral current events in these two regions, for an exposure of 1 year, are enough to obtain an error on the strange axial-charge of 6×10^{-3}, more than 10 times smaller than available at present.

PACS numbers: 25.30.Pt; neutrino scattering. 14.20.Dh; properties of protons.

Introduction: The neutrino-proton elastic scattering was proposed by Weinberg [1] as a tool to investigate the neutral currents (NC). However, this reaction depends upon 'proton strangeness'–i.e., the strange-quark contribution to the axial form-factor of the proton–a quantity that at present is not reliably known. Several theoretical [2] as well experimental investigations of neutrino-proton elastic scattering [3–5] tried to probe this quantity. The best experimental result [3] found a value for the proton strangeness similar to the one predicted in [2], but the 90% C.L. range is compatible with zero.

Proton strangeness matters in many contexts, e.g.: for the measurement of supernova neutrinos [6], for the couplings of certain dark matter candidates to nucleons [7], for the polarized parton distribution functions, e.g., [8]. However, the flavor singlet term measured by deep inelastic scattering of charged leptons includes an anomalous gluonic contribution [9]; unless one postulates that this contribution is zero [10], a considerable amount of additional labor is needed to extract proton strangeness, see e.g., [11]. By contrast, a measurement of elastic neutrino-proton scattering probes this quantity directly.

In the recent years, neutrino detectors underwent major developments. The Borexino detector, in particular, has reached an unprecedented low energy threshold for the detection of solar neutrinos [12], implying the possibility to use highly pure scintillator detectors to measure protons with kinetic energies as low as 1.3 MeV. Thus, these technological achievements pave the way for new approaches to measure precisely the elastic scattering of neutrinos onto protons.

We study the possibility to determine the proton strangeness measuring *low energy neutrino-proton scattering*. The advantages of the low energy regime are discussed in the next section. An artificial neutrino beam produced by pion decay at rest, as in the cyclotrons considered by the DAEδALUS proposal [13, 14] or in the SNS facility [15] is considered. Recall that the neutrino spectra from pion decay are very well known, and have already proved being useful for measurements of other cross sections [16]. The events are observed by a scintillating detector. We consider a 1 kton mass detector, located at 100 m from the source and with the performances of Borexino [12]. We explore the potential of this experimental setup and determine which sensitivity can be reached with 1 year of exposure. We discuss the dependence of the sensitivity on the experimental features, showing how to rescale our findings for different configurations; i.e., by varying the detector mass, the distance to the source, the time of data taking, etc..

Advantages of Low Energies: The neutral current elastic scattering transition $\nu(k)+p(p)\to\nu(k')+p(p')$ is described by the invariant matrix element

$$\mathcal{M}=-\frac{G_F}{\sqrt{2}}\;\bar{u}'_\nu\gamma^a P_L u_\nu\;\bar{u}'_p\Gamma_a(q)u_p \tag{1}$$

with

$$\Gamma_a(q)=\gamma_a F_1(Q^2)+i\frac{\sigma_{ab}q^b}{2m_p}F_2(Q^2)-\gamma_a\gamma_5 G_{\rm A}(Q^2) \tag{2}$$

and similarly for antineutrinos, where $q=p-p'$ is the transfered momentum and $Q^2=-q^2$. The vector form factors $F_{1,2}(Q^2)$ can be obtained by applying the conserved vector current hypothesis neglecting the strange vector form factors. The most important quantity for us is the axial form factor $G_{\rm A}(Q^2)$. This can be determined from the analogous term in the charged current interactions, but only up to the SU(2) isosinglet term, due to the strange quark axial current. Its effect has been described by the parameter η:

$$G_{\rm A}(Q^2)=\frac{g_{\rm A}(0)\times(1+\eta)}{2\;(1+Q^2/M_{\rm A}^2)^2} \tag{3}$$

where the dipole parameterization of the axial form factor is used and we have $g_{\rm A}(0)=1.267$. Adopting the usual

---

[*] giulia.pagliaroli@lngs.infn.it
[†] carolup@fisica.ugto.mx
[‡] manimala.mitra@lngs.infn.it
[§] francesco.vissani@lngs.infn.it



FIG. 1. *Variation of $\mathcal{R}(E_\nu)$ with the neutrino/antineutrino energy E_ν. The black solid line corresponds to $\eta = 0.12$, while in the red dashed and brown dotted line η has been set to zero. The red dashed line represents the effect of a relatively higher axial mass $M_A = 1.2$ GeV in the numerator of Eq.6. See the text for detail.*

definition $\langle p|\bar{q}\gamma^a\gamma^5 q|p\rangle \equiv \bar{u}_p\gamma^a\gamma^5 u_p \cdot \Delta q$ the electroweak standard model predicts $G_{\rm A}(0) = \Delta u/2 - \Delta d/2 - \Delta s/2$, which agrees with Eq. (3) setting $g_{\rm A}(0) \equiv \Delta u - \Delta d$ and

$$\eta \equiv -\Delta s/g_{\rm A}(0). \tag{4}$$

In this work, the term 'proton strangeness' indicates η; the conversion to Δs is evidently trivial. As recalled in the introduction, the BNL 734 experiment [3] obtained

$$\eta = 0.12 \pm 0.07 \tag{5}$$

that implies $\eta = 0 - 0.25$ at 90% C.L., compatible with zero. Moreover, due to the use of neutrinos with energies around 1 GeV, the experimental determination of η is strongly entangled with the high Q^2 behavior of the axial form factor, that is to some extent uncertain. Conversely, the impact of the 'axial mass' on the cross section is greatly reduced if lower energies are considered.

This is clear from Fig. 1, where we display the following cross section ratio:

$$\mathcal{R}(E_\nu) = \frac{\sigma_{\nu p}(E_\nu)' + \sigma_{\bar{\nu}p}(E_\nu)'}{\sigma_{\nu p}(E_\nu) + \sigma_{\bar{\nu}p}(E_\nu)}. \tag{6}$$

In the denominator of Eq. (6), we put the reference cross section, as given in [3], with $\eta = 0$ and $M_A = 1.032$ GeV. In the numerator we modified it as follows: 1) In the black curve, we use the central value $\eta = 0.12$ of Eq. (5) instead. Evidently, this causes an increase in the cross section. 2) In the red (dashed) curve, we set $\eta = 0$ and use $M_A = 1.2$ GeV. By increasing M_A, the axial form factor and thus the cross section increase. 3) In the brown (dotted) curve, we set $\eta = 0$ and omit the vector current contribution altogether. Thus, at low energies the vector form factors are practically irrelevant.

We conclude that at GeV energies, the effect of a nonzero η can be mimicked by a larger value of M_A. On the other hand, for $E <$ few 100 MeV, the cross section depends only on the $G_{\rm A}(0)$ and the interpretation of a precise measurement is much simpler, leading directly to a measurement of η. These are the reasons of interest in considering experiments with low energy neutrinos, despite the fact that the neutrino-proton cross section diminishes strongly (quadratically) with neutrino energy.

Observable Spectra in Ultra-pure Scintillators: Let us consider an intense beam of neutrinos produced from pion decay at rest according to the usual decay chains $\pi^+ \to \mu^+ + \nu_\mu \to e^+ + \nu_e + \bar{\nu}_\mu + \nu_\mu$. The energy spectra of these neutrinos are very well-known and include a monochromatic line for the muonic neutrinos at $(m_\pi^2 - m_\mu^2)/2m_\pi = 29.8$ MeV and two continuous spectra with energy between $0 < E_\nu < (m_\mu^2 - m_e^2)/2m_e = 52.8$ MeV. We assume $N_{\pi^+} = 4.9 \times 10^{22}$ pions/yr and thus, the same number of ν_e, ν_μ and $\bar{\nu}_\mu$. Such a pion production rate can be provided, for example, by a cyclotron with 800 MeV kinetic energy of protons and with a peak power of 1 MW [13].

We calculate the neutral current interactions of these ν_e, ν_μ and $\bar{\nu}_\mu$ with the protons of a ultra-pure scintillator detector of mass 1 kton located at a distance of 100 meters and with the low energy performances as of Borexino detector. The expected distribution of the elastic scattering events, $\nu(\bar{\nu}) + p \to \nu(\bar{\nu}) + p$, in the kinetic energy of the final state proton $T_p \equiv E_p' - m_p$ can be written as

$$\begin{aligned}\frac{dN}{dT_p} = N_p \int_{E_{\rm min}} dE_\nu \Bigg(&\frac{d\sigma_\nu}{dT_p}(E_\nu, T_p)\ \Phi_\nu(E_\nu) + \\ &+ \frac{d\sigma_{\bar{\nu}}}{dT_p}(E_\nu, T_p)\ \Phi_{\bar{\nu}}(E_\nu)\Bigg)\end{aligned} \tag{7}$$

where N_p is the number of protons in the scintillator, $\sigma_{\nu,\bar{\nu}}$ are the cross sections and Φ_ν and $\Phi_{\bar{\nu}}$ are the total neutrino and antineutrino fluences (i.e., time integrated fluxes) differential in the neutrino energy E_ν. The kinetic energy T_p of the proton in the final state and the minimum energy of the neutrino in the initial state are related as:

$$E_{\rm min} = \frac{1}{2}\left[T_p + \sqrt{(2m_p + T_p)T_p}\right]. \tag{8}$$

The proton kinetic energy is converted into an 'equivalent' detectable electron energy E_d, the *part* of T_p that goes into scintillation light. This can be calculated according to the empirical Birks formula, sometimes called 'quenching' function, [6, 17, 18]

$$E_d = \int_0^{T_p} \frac{dT}{1 + k_{\rm B}\,\langle dT_p/dx\rangle} \tag{9}$$

where $\langle dT_p/dx\rangle$ is the stopping power of the proton that depends on the chemical composition of the detector.

For the purpose of the analysis we use a Gaussian energy resolution $\sigma(E_d) = 50\ \text{keV}\sqrt{E_d/\text{MeV}}$ similar to Borexino [12]. We assume the chemical composition C_9H_{12}, finding that the number of protons in 1 kton is



FIG. 2. *Expected event distribution in the detectable energy* E_d *in 1 kton of scintillator as Borexino. The thick solid and dotted black lines show the event spectrum for neutral current elastic interactions of neutrinos and antineutrinos with protons. The dashed blue line represents the cosmogenic* ^{11}C *background. The magenta, pink, gray and black solid lines represent* ^{8}B*, CNO,* ^{7}Be *and PeP solar neutrino spectrum. The purple, red and green dashed lines represent the backgrounds due to* ^{210}Po*,* ^{210}Bi *and* ^{85}Kr*, respectively [21].*

$N_p = 6.02 \times 10^{31}$ (this increases only by 20% for the composition $\mathrm{C_6H_5C}_n\mathrm{H}_{2n+1}$ with $n = 12$ considered for LENA [19]). The Birks Law is obeyed at the percent level with a Birks constant of $k_\mathrm{B} = 0.0175$ [20]. Therefore, the detection energy threshold $E_d^{th} = 0.25$ MeV implies that the equivalent proton kinetic energy threshold, calculated with Eq. (9), is about 1.3 MeV. Using Eq. (8), we see that we are sensitive only to the highest part of the neutrino energy spectra, namely, $E_\nu \geq 25$ MeV.

The differential spectrum dN/dE_d expected for 1 year of exposure is shown in Fig. 2. The signal due to NC νp interactions is plotted using three black thick lines. They represent the events expected assuming $\eta = 0.0, 0.12$ (best fit of [3]) and 0.25. From Fig. 2 we see that the energy window of interest for the detection of the elastic scattering interactions is $E_d = (0.25 - 2.2)$ MeV. The number of signal events expected in this energy window and for one year of exposure is considerable

η	signal events
0.00	0.89×10^5
0.12	1.14×10^5
0.25	1.41×10^5

However, the same energy window will also contain several solar as well as radioactive background events. In particular solar neutrino events from ^{7}Be, CNO, PeP and ^{8}B are not negligible; we expect about 28.6, 3.0, 2.1 and 0.5 counts per day (cpd) in each 100 tons of scintillator respectively. Moreover, the backgrounds due to radioactive sources as ^{210}Po, ^{210}Bi and ^{85}Kr, as well as the cosmogenic ^{11}C have to be considered. Assuming the counting rate of the present Borexino configuration [21], we expect that each 100 tons of scintillator yield 10^3 cpd from ^{210}Po, 15 cpd from ^{210}Bi, and finally, 25 cpd from both ^{85}Kr and ^{11}C. The spectra for these backgrounds have also been displayed in Fig. 2.

Note incidentally that the Borexino detector is still improving on the background, and in the most recent runs, the ^{210}Po contaminant has already been halved [22]. The contribution of the ^{11}C is particularly critical for our purposes, and depends strongly on the depth of the underground site of the detector [23]. The value of this cosmogenic component is estimated to be 3 times smaller in LENA [19] and much less in SNO+ [24]. However, in the present study we will adopt the conservative background levels discussed above. Having outlined the expectations, we discuss now the sensitivity of the desired measurement.

Sensitivity to proton strangeness η: From Fig. 2, it is evident that background sources can pollute the extraction of the signal. However, following [25], we proceed to extract the signal from the total number of observed events, measuring the background events and subtracting them.

If we collect $N = S + B$ events in a time T and $N_0 = B \cdot T_0 / T$ events in a time T_0, when the signal is absent, we expect $S = N - N_0 \cdot T / T_0$ signal events. Since both N and N_0 are subject to Poissonian fluctuations, the fractional uncertainty caused by the statistical effects is given by:

$$\frac{\Delta S}{S} = \sqrt{\frac{1 + \frac{B}{S}\left(1 + \frac{T}{T_0}\right)}{S}} \tag{10}$$

If observations are binned in energy, we define $2/\Delta S^2 \equiv \partial^2(\chi^2)/\partial S^2$ using $\chi^2 = (S_i - \bar{S}_i)^2/\Delta S_i^2$ and $S_i = S\rho_i$, with $\sum_i \rho_i = 1$. In the minimum of the χ^2, $S_i = \bar{S}_i$,

$$\frac{\Delta S}{S} = \left(\sum_{i=1}^{n} \frac{S_i}{1 + \frac{B_i}{S_i}\left(1 + \frac{T}{T_0}\right)}\right)^{-1/2} \tag{11}$$

This is smaller than Eq. (10) where we set $S = \sum_i S_i$ and $B = \sum_i B_i$, unless the signal and the background have the same shape. For constant background-signal ratio, B_i/S_i =constant, both the results coincide.

From Fig. 2 it is evident that there are two comparatively clean energy windows to extract the signal, one between $(0.65 - 1.1)$ MeV and another between $(1.73 - 2.2)$ MeV respectively. Using $T = T_0$ =1 year, we expect $S_1 = 28900$ and $B_1 = 37200$ events from the first energy window and $S_2 = 1900$ and $B_2 = 2400$ events from the latter one. Since we have $B_1/S_1 \simeq B_2/S_2$ we can use Eq. (10) obtaining $\Delta S/S = 1.1$ %, dominated by the data in the first energy window. If we consider that for a typical pulsed signal [13], T_0 is 4 times larger than $T = 1$, the error decreases to $\Delta S/S = 0.9\%$; a similar improvement is obtained if we measure the background over the whole range $0.65 < E_d < 2.5$ MeV, increasing the statistic and thus reducing its error.

The NC cross section σ at low energy (and therefore the signal event rate S) is dominated by the axial form factor, scaling as $(1+\eta)^2$, as is clear from Eq. (3). So we can relate the signal sensitivity to the one on η through $\delta S = \Delta\sigma/\sigma = 2\Delta\eta/(1+\eta)$. This means that the sensitivity $\Delta S/S = 1.1\%$ on the NC signal implies that proton strangeness can be measured with an absolute error of $\Delta\eta = 6 \cdot 10^{-3}$ for $\eta = 0.12$, i.e., a 5% fractional uncertainty. For all values of η in the allowed experimental range, the error is more than 10 times smaller than Eq. (5).

Finally, we show how to attain the same sensitivity with different experimental parameters. Let us write the uncertainty in η as follows,

$$\Delta\eta = \frac{1+\eta}{2}\sqrt{\frac{1+\frac{1}{\varphi}\frac{B}{S}\left(1+\tau\,\frac{T}{T_0}\right)}{\varepsilon\,\varphi\,S}} \qquad (12)$$

Here $\varepsilon = (M'/M)(T'/T)$ rescales the exposure, proportional to the mass M' and to the time of data taking T'; $\tau = (T'/T_0')/(T/T_0)$ rescales the fraction of data taking time that includes the signal; $\varphi = (P'/P)(D/D')^2$ rescales the flux, accounting for the new power P' and the new distance D'. Let us adopt the operational parameters of [13], i.e., assume a distance from the cyclotron of $D' = 700$ m, a power increased by a factor of 5 and $T' = T_0'/4 = 1$ year. The flux (and the signal) decreases by one order of magnitude, for it increases with the power but decreases as the distance squared. In order to compensate for this, we need a larger exposure. E.g., with a mass of 50 kton as in the LENA proposal [19] we obtain again $\Delta S/S = 1.1$ % after 1 year. Note that the selection of the optimal distance between source and detector will require to minimize the beam related backgrounds.

Summary: In this work, we explored the possibility to study proton strangeness using low energy neutrinos. We discussed the potential of exploiting a synergy between artificial neutrino beams from pion decay and ultra-pure scintillating detectors. Incidentally, such complex will allow us to quantify precisely the response of underground neutrino detectors, to search for sterile neutrinos and to investigate many more issues.

We have shown that, by using reasonable assumptions on the experimental parameters, it is possible to identify two energy windows that allow to measure many tens of thousand of signal events. They imply a statistical error on proton strangeness one order of magnitude smaller than obtained by BNL 734, Eq. (5). It is important to emphasize that the signal we discussed should be collected above 0.6 MeV, and thus, it does not require the extreme low energy threshold reached by Borexino detector.

ACKNOWLEDGMENTS

We thank warmly A. Bacchetta, G. Bruno, G.T. Garvey, W. Fulgione, A. Ianni, M. Mannarelli, A. Molinario and F. Villante for help and useful discussions.

[1] S. Weinberg, Phys. Rev. D **5** (1972) 1412.

[2] J. C. Collins, F. Wilczek and A. Zee, Phys. Rev. D **18** (1978) 242. R. Mohapatra and G. Senjanovic, Phys. Rev. D **19** (1979) 2165. L. Wolfenstein, Phys. Rev. D **19** (1979) 3450.

[3] L.A. Ahrens *et al.* [BNL 734] Phys. Rev. D **35** (1987) 785.

[4] R. Tayloe, Nucl.Phys.B (Proc. Suppl.) **105** (2002) 62.

[5] A.A. Aguilar-Arevalo *et al.* [MiniBooNE] Phys. Rev. D **82** (2010) 092005.

[6] J. F. Beacom, W. M. Farr and P. Vogel, Phys. Rev. D **66** (2002) 033001.

[7] J. Ellis and M. Karliner, Phys. Lett. B **341** (1995) 397.

[8] J. Ashman *et al.* [EMC], Nucl. Phys. B **328** (1989) 1.

[9] A.V. Efremov and O.V. Teryaev, JINR Report E2-88-287, Dubna (1988); G. Altarelli and G.G. Ross, Phys. Lett. B 212 (1988) 391.

[10] W. M. Alberico, S. M. Bilenky and C. Maieron, Phys. Rept. **358** (2002) 227.

[11] D. de Florian, R. Sassot, M. Stratmann and W. Vogelsang, Phys. Rev. Lett. **101** (2008) 072001.

[12] C. Arpesella *et al.* [Borexino], Phys. Rev. Lett. **101** (2008) 091302.

[13] J. Alonso, F. T. Avignone, W. A. Barletta, R. Barlow, H. T. Baumgartner, A. Bernstein, E. Blucher and L. Bugel *et al.*, arXiv:1006.0260 [physics.ins-det].

[14] A. Bungau *et al.* [DAEδALUS and for the DAEδALUS Collaborations], arXiv:1205.5528 [physics.ins-det].

[15] M. Sajjad Athar, S. Ahmad and S. K. Singh, Nucl. Phys. A **764** (2006) 551.

[16] B. Bodmann *et al.* [KARMEN] Phys. Lett. B **267** (1991) 321.

[17] J. B. Birks, Proc. of the Phys. Soc. A **64** (1951) 874.

[18] W. R. Leo, *Techniques for Nuclear and Particle Physics Experiments* Second Edition (Springer-Verlag, Berlin, 1994).

[19] M. Wurm *et al.* [LENA], Astropart. Phys. **35** (2012) 685.

[20] A. Ianni, Proc. of the HAνSE 2011 workshop, Hamburg, July 19-23, 2011, ed. A. Mirizzi, P. Serpico, G. Sigl, Verlag Deutsches Elektronen-Sychrotron.

[21] G. Bellini *et al.* [Borexino], Phys. Rev. Lett. **107** (2011) 141302. G. Bellini *et al.* [Borexino], Phys. Rev. Lett. **108** (2012) 051302.

[22] A. Ianni, talk given at NOW 2012, available at `http://www.ba.infn.it/~now/`.

[23] C. Galbiati *et al.*, Phys. Rev. C **71** (2005) 055805.

[24] C. Kraus *et al.* [SNO+], Prog. Part. Nucl. Phys. **64** (2010) 273.

[25] A. Ianni, D. Montanino and F. L. Villante, Phys. Lett. B **627** (2005) 38.